

cdf

DIRECTIONS FOR USE Mix well before use.

Cotton

To improve production, Phytamin® All Purpose should be applied as a foliar spray during the early bloom

Application Rates: ½ - 2 gallons per acre in sufficient water to assure thorough coverage. Begin treatments at first bloom up to 3 weeks after first bloom. Treatments may be made up to three times at 10 day intervals,

Trees, Fruit & Nuts

Application Rate: 1/2 - 1 gallon per acre applied in ample water to provide for thorough coverage. Begin treatments between first bloom and full bloom. Repeat

Grapes

Application Rate: ½ - 1 gallons per acre applied at 7 to 14 day intervals starting when spring cane growth reaches 12 inches in length.

Tomatoes

Phytamin® All Purpose may be applied at any time throughout the bloom period.

Application Rate: 1/2 - 1 gallon per acre at 10 to 14 day intervals throughout the bloom period.

Peppers

Phytamin® All/Purpose should be applied during the

Application Rate: 1/2 - 1 gallon per acre applied in sufficient water for thorough coverage. Two to three treatments should be made at 10 day intervals

Vegetables

Do not apply as a foliar fertilizer to lettuce or other

Yield and quality may be improved by regular treatments of Phytamin® All Purpose throughout the

Application Rates: 1/2 - 1 gallon per acre in sufficient water for thorough coverage. Treatments should be made after thinning at 10 to 14 day intervals.

Other Crops

Application Rates: Phytamin® All Purpose may be used successfully on most crops. Please consult your dealer for rates and timing of applications.

SOIL APPLICATION

Side-dress: Apply 5 to 50 gallons per acre. Drip Systems: Apply 1 to 10 gallons per acre per week. Water-run: Apply 3 to 30 gallons per acre. Pre-plant: Apply 2 to 50 gallons per acre. Sprinkler Irrigation: Apply 2 to 20 gallons per acre.

🗌 275 gal. / 1041 L

Bulk

Density: 10 lbs. per gallon at 68° F